arXiv:1704.07149v1 [cs.LO] 24 Apr 2017

Axiomatizing Epistemic Logic of Friendship via Tree Sequent Calculus

Katsuhiko Sano

Department of Philosophy, Graduate School of Letters, Hokkaido University, Japan
v-sano@let.hokudai.ac.jp

Abstract. This paper positively solves an open problem if it is possible to provide a Hilbert system to Epistemic Logic of Friendship (EFL) by Seligman, Girard and Liu. To find a Hilbert system, we first introduce a sound, complete and cut-free tree (or nested) sequent calculus for EFL, which is an integrated combination of Seligman's sequent calculus for basic hybrid logic and a tree sequent calculus for modal logic. Then we translate a tree sequent into an ordinary formula to specify a Hilbert system of EFL and finally show that our Hilbert system is sound and complete for the intended two-dimensional semantics.

Keywords: Epistemic Logics of Friendship, Tree Sequent Calculus, Hilbert System, Completeness, Cut Elimination Theorem

1 Introduction

Epistemic Logic of Friendship (**EFL**) is a version of two-dimensional modal logic proposed by [20,19]. Compared to the ordinary epistemic logic [12], one of the key features of their logic is to encode the information of agents into the object language by a technique of hybrid logic [3,1]. Then, a propositional variable p can be read as an indexical proposition such as 'I am p' and we may formalize the sentences like 'I know that all of my friend is p' or 'All of my friends know that he/she is p'. Moreover, they provide a dynamic mechanism for capturing public announcements [16], announcements to all the friends, and private announcements [2] and established a relative completeness result (cf. [20,10]). This paper focuses on the problem of axiomatizing **EFL** in terms of Hilbert system, i.e., the static part of their framework.

A difficulty of the problem comes from a combination of *modal logic* for agents' knowledge and *hybrid logic* for a friendship relation among agents. If we combine *two hybrid logics* over two-dimensional semantics of [20,19], it is noted that there is an axiomatization of all valid formulas in the semantics by [17, p.471]. Our approach to tackle the problem is via a sequent calculus, whose idea is originally from Gentzen. In particular, our notion of sequent for **EFL** can be regarded as a combination of a tree or nested sequent [13,6] for modal logic and @-prefixed sequent [18,5] for hybrid logic. One of the merits of our notion of sequent is that we can translate our sequent into an ordinary formula. This allows us to specify our desired Hilbert system for **EFL**. We note that [7] independently provided a prefixed tableau system for a dynamic extension of **EFL**. There are at least three points we should emphasize on our work. First, our tree

sequent system is quite simpler than the tableau system given in [7], i.e., the number of rules of our sequent system is almost half of the number of rules of their system. Second, it is not clear if a prefixed formula in [7] for the tableau calculus can be translated into an ordinary formula. Their result is not concerned with Hilbert system. Third, their syntax contains a special kind of propositional variable (called *feature proposition*) and they include a tableau rule called *propositional cut* to handle such propositions. On the other hand, we can show that our tree sequent calculus enjoys the cut elimination theorem, the most fundamental theorem in proof-theory.

We proceed as follows. Section 2 introduces the syntax and semantics of $\mathbf{EFL}$. Section 3 provides a tree sequent calculus for $\mathbf{EFL}$ and establish the soundness of the sequent calculus (Theorem 1). Section 4 establish a completeness result of a cut-free fragment of our sequent calculus (Theorem 2). As a corollary, we also provide a semantic proof of the cut elimination theorem of our sequent calculus (Corollary 1). Section 5 specifies a Hilbert system of $\mathbf{EFL}$, and provide a syntactic proof of the equipollence between our proposed Hilbert system and our tree sequent calculus, which implies the soundness and completeness results for our Hilbert system (Corollary 2). Section 6 extends our technical results to cover extensions of $\mathbf{EFL}$ where $\Box$ (a modal operator for states, or a knowledge operator) obeys $\mathbf{S4}$ or $\mathbf{S5}$ axioms and a friendship relation satisfies some universal properties (Theorem 5). The result of this section subsumes the logic given in [7], provided we drop the dynamic operator from the syntax of [7]. Section 7 concludes this paper. [1]

2 Syntax and Two-dimensional Kripke Semantics

Our syntax $\mathcal{L}$ consists of the following vocabulary: a countably infinite set $\mathsf{Prop} = \{ p, q, r, \dots \}$ of propositional variables, a countably infinite set $\mathsf{Nom} = \{ n, m, l, \dots \}$ of agent nominal variables, the Boolean connectives of $\to$ (the implication) and $\bot$ (the falsum), the satisfaction operators $@$ and the friendship operator F as well as the modal operator $\Box$. We note that an *agent nominal* $n \in \mathsf{Nom}$ is a syntactic name of an agent or an individual, which amounts to a constant symbol of the first-order logic, while n is read indexically as "I am n." Similarly, we read a propositional variable $p \in \mathsf{Prop}$ also indexically by "I am p," e.g., "I am in danger." The set Form of formulas in $\mathcal{L}$ is defined inductively as follows:

$$\mathsf{Form} \ni \varphi ::= n \mid p \mid \bot \mid \varphi \to \varphi \mid @_n \varphi \mid \mathsf{F}\varphi \mid \Box\varphi,$$

where $n \in \mathsf{Nom}$ and $p \in \mathsf{Prop}$. Boolean connectives other than $\to$ or $\bot$ are introduced as ordinary abbreviations. We define the dual of $\Box$ as $\Diamond := \neg\Box\neg$ and the dual of F as $\langle \mathsf{F} \rangle := \neg \mathsf{F} \neg$. Moreover, a formula of the form $@_n\varphi$ is said to to be $@$-*prefixed*. Let us read $\Box$ as "I know that." Here are some examples of how to read formulas:

[1] I presented the contents of this paper first at the 48th MLG meeting at Kaga, Ishikawa, Japan on 6th December 2013 and then at Kanazawa Workshop for Epistemic Logic and its Dynamic Extensions, Kanazawa, Japan on 22nd February 2014. I would like to thank Jeremy Seligman and Fenrong Liu for fruitful discussions. All errors, however, are mine. The work of the author was partially supported by JSPS KAKENHI Grant-in-Aid for Young Scientists (B) Grant Number 15K21025 and JSPS Core-to-Core Program (A. Advanced Research Networks).

- $\Box p$, read as "I know that I am p."
- $@_n \Box p$, read as "n knows that she is p."
- $\Box @_n p$, read as "I know that agent n is p."
- $\mathsf{F}p$, read as "all of my friends are p."
- $\mathsf{F}\Box p$, read as "all of my friends know that they are p."
- $\Box \mathsf{F} p$, read as "I know that all of my friends are p."
- $@_n \langle \mathsf{F} \rangle m$, read as "agent m is a friend of agent n."

We say that a mapping $\sigma : \mathsf{Prop} \cup \mathsf{Nom} \to \mathsf{Form}$ is a *uniform substitution* if σ uniformly substitutes propositional variables by formulas and agent nominals by agent nominals and we use $\varphi\sigma$ to mean the result of applying a uniform substitution σ to φ. In particular, we use $\varphi[n/k]$ to mean the result of substituting each occurrence of agent nominal k in φ uniformly with agent nominal n.

An *model* $\mathfrak{M}$ for our syntax $\mathcal{L}$ is a tuple $(W, A, (R_a)_{a \in A}, (\asymp_w)_{w \in W}, V)$, where W is a non-empty set of possible states, A is a non-empty set of agents, R_a is a binary relation on W ($a \in A$), $\asymp_w$ is a binary relation on A (called a *friendship relation*), V is a valuation function $\mathsf{Prop} \cup \mathsf{Nom} \to \mathcal{P}(W \times A)$ such that $V(n)$ is a subset of $W \times A$ of the form $W \times \{ a \}$, where we denote such unique element a by $\underline{n}$. We do not require any property for R_a and $\asymp_w$ but we will come back to this point in Section 6. We say that a tuple $\mathfrak{F} = (W, A, (R_a)_{a \in A}, (\asymp_w)_{w \in W})$ without a valuation is a *frame*.

Let $\mathfrak{M} = (W, A, (R_a)_{a \in A}, (\asymp_w)_{w \in W}, V)$ be a model. Given a pair $(w, a) \in W \times A$ and a formula φ, the satisfaction relation $\mathfrak{M}, (w, a) \models \varphi$ (read "agent a satisfies φ at w in $\mathfrak{M}$ ") inductively as follows:

$$
\begin{array}{ll}
\mathfrak{M}, (w,a) \models p & \text{iff } (w,a) \in V(p), \\
\mathfrak{M}, (w,a) \models n & \text{iff } \underline{n} = a, \\
\mathfrak{M}, (w,a) \not\models \bot & \\
\mathfrak{M}, (w,a) \models \varphi \to \psi & \text{iff } \mathfrak{M}, (w,a) \models \varphi \text{ implies } \mathfrak{M}, (w,a) \models \psi \\
\mathfrak{M}, (w,a) \models @_n \varphi & \text{iff } \mathfrak{M}, (w, \underline{n}) \models \varphi, \\
\mathfrak{M}, (w,a) \models \mathsf{F}\varphi & \text{iff } (a \asymp_w b \text{ implies } \mathfrak{M}, (w,b) \models_{\mathfrak{M}} \varphi) \text{ for all agents } b \in A, \\
\mathfrak{M}, (w,a) \models \Box \varphi & \text{iff } (w R_a v \text{ implies } \mathfrak{M}, (v,a) \models \varphi) \text{ for all states } v \in W.
\end{array}
$$

Given a class $\mathbb{M}$ of models, we say that a formula φ is *valid* in $\mathbb{M}$ when $\mathfrak{M}, (w, a) \models \varphi$ for all pairs (w, a) in $\mathfrak{M}$ and all models $\mathfrak{M} \in \mathbb{M}$. This paper tackles the question if the set of all valid formulas in the class of all models is axiomatizable.

3 Tree Sequent Calculus of Epistemic Logic of Friendship

A *label* is inductively defined as follows: Any natural number is a label; if α is a label, n is an agent nominal in Nom and i is a natural number, then $\alpha \cdot_n i$ is also a label. When β is $\alpha \cdot_n i$, then we say that β is an *n-child* of α or that α is *an n-parent* of β. A *tree* $\mathcal{T}$ is a set of labels such that the set is closed under taking the parent of a label, i.e., $\alpha \cdot_n i \in \mathcal{T}$ implies $\alpha \in \mathcal{T}$ for all labels α, agent nominals n and natural numbers i. For example, all of 0, $0 \cdot_n 1$ and $0 \cdot_k 2$ are labels and they form a tree.



Fig. 1. A tree sequent

Given a label α and an $@$-*prefixed formula* φ, the expression $\alpha : \varphi$ is said to be a *labelled formula*, where recall that an $@$-prefixed formula is of the form $@_n\varphi$. A *tree sequent* is an expression of the form

$$\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$$

where Γ and Δ are *finite* sets of labelled formulas, $\mathcal{T}$ is a tree of labels, and all the labels in Γ and Δ are in $\mathcal{T}$. A tree sequent "$\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$" is read as "if we assume all labelled formulas in Γ, then we may conclude some labelled formulas of Δ." A tree sequent $0 : @_n\varphi, 0 \cdot_k 2 : @_m\rho \overset{\mathcal{T}}{\Rightarrow} 0 : @_m\psi, 0 \cdot_n 1 : @_k\theta$ is represented as in Fig 1, where $\mathcal{T} = \{ 0, 0 \cdot_n 1, 0 \cdot_k 2 \}$. That is, 0, $0 \cdot_n 1$ and $0 \cdot_k 2$ are "addresses" of the root, the right leaf, and the left leaf, respectively.

Table 1. Tree Sequent Calculus $\mathsf{T}\mathbf{EFL}$

$$(\bot)\quad \alpha : @_n\bot, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta \qquad (\mathsf{id})\quad \alpha : @_n\varphi, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta, \alpha : @_n\varphi$$

$$\frac{\alpha : @_n m, \alpha : \varphi[n/k], \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}{\alpha : @_n m, \alpha : \varphi[m/k], \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}\ (\mathsf{rep}_{=1}) \qquad \frac{\alpha : @_n m, \alpha : \varphi[m/k], \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}{\alpha : @_n m, \alpha : \varphi[n/k], \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}\ (\mathsf{rep}_{=2})$$

$$\frac{\alpha : @_n n, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}\ (\mathsf{ref}_{=}) \qquad \frac{\beta : @_n m, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}{\alpha : @_n m, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}\ (\mathsf{rigid}_{=})$$

$$\frac{\alpha : @_n\varphi, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta, \alpha : @_n\psi}{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta, \alpha : @_n(\varphi \to \psi)}\ (\to R) \qquad \frac{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta, \alpha : @_n\varphi \quad \alpha : @_n\psi, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}{\alpha : @_n(\varphi \to \psi), \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}\ (\to L)$$

$$\frac{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta, \alpha : @_m\varphi}{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta, \alpha : @_n@_m\varphi}\ (@R) \qquad \frac{\alpha : @_m\varphi, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}{\alpha : @_n@_m\varphi, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}\ (@L)$$

$$\frac{\alpha : @_n\langle\mathsf{F}\rangle m, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta, \alpha : @_m\varphi}{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta, \alpha : @_n\mathsf{F}\varphi}\ (\mathsf{F}R)^* \qquad \frac{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta, \alpha : @_n\langle\mathsf{F}\rangle m \quad \alpha : @_m\varphi, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}{\alpha : @_n\mathsf{F}\varphi, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}\ (\mathsf{F}L)$$

$$\frac{\Gamma \overset{\mathcal{T}\cup\{\alpha\cdot_n i\}}{\Rightarrow} \Delta, \alpha \cdot_n i : @_n\varphi}{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta, \alpha : @_n\Box\varphi}\ (\Box R)^\dagger \qquad \frac{\beta : @_n\varphi, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}{\alpha : @_n\Box\varphi, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}\ (\Box L)^\ddagger$$

$$\frac{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}{\Gamma \overset{\mathcal{T}\cup\{\alpha\}}{\Rightarrow} \Delta}\ (w\mathsf{lab})^\star \qquad \frac{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta, \alpha : @_n\varphi \quad \alpha : @_n\varphi, \Pi \overset{\mathcal{T}}{\Rightarrow} \Sigma}{\Gamma, \Pi \overset{\mathcal{T}}{\Rightarrow} \Delta, \Sigma}\ (Cut)$$

$*$: m is a fresh agent nominal in the lower sequent; $\dagger$: $i \in \mathbb{N}$ is fresh in the lower sequent;
$\ddagger$: β is an n-child of α; $\star$: $\mathcal{T} \cup \{\alpha\}$ is a tree of labels

Table 1 provides all the initial sequents and all the inference rules of tree sequent calculus $\mathsf{T}\mathbf{EFL}$, where $\Gamma[m/k]$ is defined as $\{ \alpha : \varphi[m/k] \mid \alpha : \varphi \in \Gamma \}$ and recall that $\varphi[m/k]$ is the result of substituting each occurrence of agent nominal k in φ with agent nominal m. The system without the cut rule is denoted by $\mathsf{T}\mathbf{EFL}^-$. A *derivation* in $\mathsf{T}\mathbf{EFL}$ (or $\mathsf{T}\mathbf{EFL}^-$) is a finite tree generated from initial sequents by inference rules of $\mathsf{T}\mathbf{EFL}$ (or $\mathsf{T}\mathbf{EFL}^-$, respectively). The *height* of a derivation is defined as the maximum length of branches in the derivation from the end (or root) sequent to an initial sequent. A tree sequent $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ is said to be *provable* in $\mathsf{T}\mathbf{EFL}$ (or $\mathsf{T}\mathbf{EFL}^-$) if there is a derivation in $\mathsf{T}\mathbf{EFL}$ (or $\mathsf{T}\mathbf{EFL}^-$, respectively) such that the root of the tree is $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$.

Let $\mathfrak{M} = (W, A, (R_a)_{a\in A}, (\succeq_w)_{w\in W}, V)$ be a model and $\mathcal{T}$ a tree of labels. A function $f : \mathcal{T} \to W$ is a *$\mathcal{T}$-assignment* in $\mathfrak{M}$ if, whenever β is an n-child of α in $\mathcal{T}$, $f(\alpha) \succeq_{\underline{n}} f(\beta)$ holds. Given any labelled formula $\alpha : @_n\varphi$ such that $\alpha \in \mathcal{T}$ and a $\mathcal{T}$-assignment in $\mathfrak{M}$, we define the *satisfaction* for a labelled formula as follows:

$$\mathfrak{M}, f \models \alpha : @_n\varphi \text{ iff } \mathfrak{M}, (f(\alpha), \underline{n}) \models \varphi.$$

where "$\mathfrak{M}, f \models \alpha : @_n\varphi$" is read as "$\alpha : @_n\varphi$ is true at $(\mathfrak{M}, f)$". Given a tree sequent $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ and a $\mathcal{T}$-assignment in $\mathfrak{M}$, $\mathfrak{M}, f \models \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ ($read$:"$\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ is true in $(\mathfrak{M}, f)$") if, whenever all labelled formulas of Γ is true in $(\mathfrak{M}, f)$, some labelled formulas of Δ is true in $(\mathfrak{M}, f)$. The following theorem is easy to establish.

Theorem 1 (Soundness of $\mathsf{T}\mathbf{EFL}$). *If a tree sequent $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ is provable in $\mathsf{T}\mathbf{EFL}$ then $\mathfrak{M}, f \models \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ for all models $\mathfrak{M}$ and all assignments f.*

Let us say that an inference rule is *height-preserving admissible* in $\mathsf{T}\mathbf{EFL}^-$ (or $\mathsf{T}\mathbf{EFL}$) if, whenever all uppersequents (premises) of the inference rule is provable by a derivation whose height is n, then the lowersequent (conclusion) of the rule is provable by a derivation whose height is at most n. By induction on height n of a derivation, we can prove the following.

Proposition 1. *The following weakening rules* ($\mathsf{w}R$) *and* ($\mathsf{w}L$) *are height-preserving admissible in $\mathsf{T}\mathbf{EFL}^-$ and $\mathsf{T}\mathbf{EFL}$. Moreover, the following substitution rule* (sub) *is height-preserving admissible in $\mathsf{T}\mathbf{EFL}^-$ and $\mathsf{T}\mathbf{EFL}$:*

$$\frac{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta, \alpha : @_n\varphi}\ (\mathsf{w}R) \qquad \frac{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}{\alpha : @_n\varphi, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}\ (\mathsf{w}L) \qquad \frac{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}{\Gamma\sigma \overset{\mathcal{T}}{\Rightarrow} \Delta\sigma}\ (\mathsf{sub}),$$

where σ is a uniform substitution and $\Theta\sigma := \{\, \alpha : \varphi\sigma \in | \, \alpha : \varphi \in \Theta \,\}$.

4 Semantic Completeness of Tree Sequent Calculus of Epistemic Logic of Friendship

In what follows in this section, multisets Γ, Δ, etc. of labelled formulas and a tree $\mathcal{T}$ of labels can be possibly (countably) infinite. Following this change, we say that a possibly infinite tree-sequent $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ is provable in $\mathsf{T}\mathbf{EFL}^-$ if there exist finite multisets $\Gamma' \subseteq \Gamma$ and $\Delta' \subseteq \Delta$ and finite subtree $\mathcal{T}'$ of $\mathcal{T}$ such that $\Gamma' \overset{\mathcal{T}'}{\Rightarrow} \Delta'$ is provable in $\mathsf{T}\mathbf{EFL}^-$.

Definition 1 (saturated tree sequent). *A possibly infinite tree sequent $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ is* saturated *if it satisfies the following conditions:*

(rep1) *If $\alpha : @_n m \in \Gamma$ and $\alpha : \varphi[n/k] \in \Gamma$ then $\alpha : \varphi[m/k] \in \Gamma$.*
(rep2) *If $\alpha : @_m n \in \Gamma$ and $\alpha : \varphi[n/k] \in \Gamma$ then $\alpha : \varphi[m/k] \in \Gamma$.*
(ref$_=$) *$\alpha : @_n n \in \Gamma$ for all labels $\alpha \in \mathcal{T}$.*
(rigid$_=$) *If $\alpha : @_n m \in \Gamma$ then $\beta : @_n m \in \Gamma$ for all labels $\beta \in \mathcal{T}$.*

($\to$r) *If* $\alpha : @_n(\varphi \to \psi) \in \Delta$ *then* $\alpha : @_n\varphi \in \Gamma$ *and* $\alpha : @_n\psi \in \Delta$.
($\to$l) *If* $\alpha : @_n(\varphi \to \psi) \in \Gamma$ *then* $\alpha : @_n\varphi \in \Delta$ *or* $\alpha : @_n\psi \in \Gamma$.
(@r) *If* $\alpha : @_n@_m\varphi \in \Delta$ *then* $\alpha : @_m\varphi \in \Delta$.
(@l) *If* $\alpha : @_n@_m\varphi \in \Gamma$ *then* $\alpha : @_m\varphi \in \Gamma$.
(Fr) *If* $\alpha : @_n\mathsf{F}\varphi \in \Delta$ *then* $\alpha : @_n\langle\mathsf{F}\rangle m \in \Gamma$ *and* $\alpha : @_m\varphi \in \Delta$ *for some nominal* m.
(Fl) *If* $\alpha : @_n\mathsf{F}\varphi \in \Gamma$ *then* $\alpha : @_n\langle\mathsf{F}\rangle m \in \Delta$ *or* $\alpha : @_m\varphi \in \Gamma$ *for all nominals* m.
($\Box$r) *If* $\alpha : @_n\Box\varphi \in \Delta$ *then* $\beta : @_n\varphi \in \Delta$ *for some* n*-child* β *of* α.
($\Box$l) *If* $\alpha : @_n\Box\varphi \in \Gamma$ *then* $\beta : @_n\varphi \in \Gamma$ *for all* n*-children* β *of* α.

By the standard argument, we can show the following *saturation lemma*.

Lemma 1. *Let* $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ *be an unprovable tree sequent in* $\mathsf{T}\mathbf{EFL}$. *Then, there exists a saturated (possibly infinite) sequent* $\Gamma^+ \overset{\mathcal{T}^+}{\Rightarrow} \Delta^+$ *such that it is still unprovable in* $\mathsf{T}\mathbf{EFL}$ *and it extends the original tree sequent, i.e.,* $\Gamma \subseteq \Gamma^+$, $\Delta \subseteq \Delta^+$ *and* $\mathcal{T} \subseteq \mathcal{T}^+$.

Lemma 2. *Let* $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ *be an saturated and unprovable tree sequent in* $\mathsf{T}\mathbf{EFL}$. *Define the derived model* $\mathfrak{M} = (\mathcal{T}, A, (R_a)_{a\in A}, (\asymp_\alpha)_{\alpha\in\mathcal{T}}, V)$ *from* $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ *by:*

- $A := \{\, |n| \mid$ n *is an agent nominal* $\}$, *where* $|n|$ *is an equivalence class of an equivalence relation* $\sim$ *which is defined as:* $n \sim m$ iff $\alpha : @_n m \in \Gamma$ *for some* $\alpha \in \mathcal{T}$.
- $\alpha R_{|n|}\beta$ *iff* β *is an* m*-child of* α *for some* $m \in |n|$.
- $|n| \asymp_\alpha |m|$ *iff* $\alpha : @_n\langle\mathsf{F}\rangle m \in \Gamma$.
- $(\alpha, |n|) \in V(m)$ *iff* $\alpha : @_n m \in \Gamma$ $(m \in \mathsf{Nom})$.
- $(\alpha, |n|) \in V(p)$ *iff* $\alpha : @_n p \in \Gamma$ $(p \in \mathsf{Prop})$.

Then, $\mathfrak{M}$ *is a model. Moreover, for every labelled formula* $\alpha : @_n\varphi$, *we have*

(i) *If* $\alpha : @_n\varphi \in \Gamma$ *then* $\mathfrak{M}, (\alpha, |n|) \models \varphi$; (ii) *If* $\alpha : @_n\varphi \in \Delta$ *then* $\mathfrak{M}, (\alpha, |n|) \not\models \varphi$.

Proof. First, let us check that $\mathfrak{M}$ is a model. First of all, note that we can easily verify that $\sim$ is an equivalence relation by the conditions $(\mathbf{ref}_=)$, $(\mathbf{rep}_i)$ and $(\mathbf{rigid}_=)$ of Definition 1. We can also check that if $n \sim m$ then $R_{|n|} = R_{|m|}$ and that if $n \sim n'$ and $m \sim m'$ then $\alpha : @_n\langle\mathsf{F}\rangle m \in \Gamma$ iff $\alpha : @_{n'}\langle\mathsf{F}\rangle m' \in \Gamma$. So both of $R_{|n|}$ and $\asymp_\alpha$ are well-defined. As for the valuation of propositional variables, when $n \sim m$ holds, the equivalence between $\alpha : @_n p \in \Gamma$ and $\alpha : @_m p \in \Gamma$ holds by the saturation conditions $(\mathbf{rep}_1)$ and $(\mathbf{rep}_2)$. For the valuation for agent nominals m, we need to check that $\{\, (\alpha, |n|) \mid \alpha : @_n m \in \Gamma \,\}$ is $\mathcal{T} \times \{\, |m| \,\}$. But this is clear from the saturation condition $(\mathbf{rigid}_=)$ and the fact that $\sim$ is an equivalence relation.

Now we move to check items (i) and (ii) by induction on φ. We only check the cases where φ is of the form $\Box\varphi$, since the other cases are easy to establish by the corresponding saturation conditions of Definition 1. Let φ be of the form $\Box\varphi$. To show (i), assume that $\alpha : @_n\Box\varphi \in \Gamma$. We need to show $\mathfrak{M}, (\alpha, |n|) \models \Box\varphi$, so let us fix any label β such that $\alpha R_{|n|}\beta$. Our goal is to show $\mathfrak{M}, (\beta, |n|) \models \varphi$. By $\alpha R_{|n|}\beta$, we can find an agent nominal $m \in |n|$ such that β is an m-child of α. It follows from $m \in |n|$ that $\gamma : @_n m \in \Gamma$ for some label γ. By $\alpha : @_n\Box\varphi \in \Gamma$ and $\gamma : @_n m \in \Gamma$, the saturation condition $(\mathbf{rep}_1)$ implies that $\alpha : @_m\Box\varphi \in \Gamma$. By the saturation condition ($\Box$l) and the fact that β is an m-child of α, we obtain $\alpha : @_m\Box\varphi \in \Gamma$. By induction hypothesis, $\mathfrak{M}, (\beta, |m|) \models \varphi$ hence we obtain our goal by $|m| = |n|$. This finishes to show (i).

For (ii), assume that $\alpha : @_n\Box\varphi \in \Delta$. By the saturation condition ($\Box\mathbf{r}$), $\beta : @_n\varphi \in \Delta$ for some n-child β of α, i.e., $\alpha R_{|n|}\beta$. By induction hypothesis, $\mathfrak{M}, (\beta, |n|) \not\models \varphi$. So we conclude that $\mathfrak{M}, (\alpha, |n|) \not\models \Box\varphi$. □

Theorem 2 (Completeness of cut-free $\mathsf{T}\mathbf{EFL}^-$). *If a tree sequent $\mathfrak{M}, f \models \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ for all models $\mathfrak{M}$ and all assignments f, then $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ is provable in $\mathsf{T}\mathbf{EFL}^-$.*

Proof. Suppose for contradiction that $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ is unprovable in $\mathsf{T}\mathbf{EFL}^-$. By Lemma 1, we can extend this tree sequent into a saturated (possibly infinite) tree sequent $\Gamma^+ \overset{\mathcal{T}^+}{\Rightarrow} \Delta^+$ which is still unprovable in $\mathsf{T}\mathbf{EFL}^-$. Let $\mathfrak{M}$ be the derived model from $\Gamma^+ \overset{\mathcal{T}^+}{\Rightarrow} \Delta^+$. Let us define $f : \mathcal{T} \to \mathcal{T}$ as the identity mapping. Then it follows from Lemma 2 that $\mathfrak{M}, f \not\models \Gamma \Rightarrow \Delta$, as required. □

By Theorem 1 and Theorem 2, the cut elimination theorem of $\mathsf{T}\mathbf{EFL}$ follows.

Corollary 1. *The following are all equivalent:*

1. $\mathfrak{M}, f \models \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ *for all models* $\mathfrak{M}$ *and all assignments* f.
2. $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ *is provable in* $\mathsf{T}\mathbf{EFL}^-$.
3. $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ *is provable in* $\mathsf{T}\mathbf{EFL}$.

Therefore, $\mathsf{T}\mathbf{EFL}$ *enjoys the cut-elimination theorem.*

5 Hilbert System of Epistemic Logic of Friendship

This section provides a Hilbert system of the epistemic logic of friendship by "translating" a tree sequent into a formula in $\mathcal{L}$. First of all, let us introduce the notion of *necessity form*, originally proposed in [11] by Goldblatt and used also in [4,9]. Necessity forms are employed to formulate an inference rule of our Hilbert system.

Definition 2 (necessity form). *Fix an arbitrary symbol $\#$ not occurring in the syntax $\mathcal{L}$. A* necessity form *is defined inductively as follows:* (i) *$\#$ is a necessity form;* (ii) *If L is a necessity form and φ is a formula, then $\varphi \to L$ is also a necessity form;* (iii) *If L is a necessity form and n is an agent nominal, then $@_n\Box L$ is also a necessity form. Given a necessity form $L(\#)$ and a formula φ of $\mathcal{L}$, we use $L(\varphi)$ to denote the formula obtained by replacing the unique occurrence of $\#$ in L by the formula φ.*

When $L(\#)$ is a necessity form of $\psi_0 \to @_n\Box(\psi_1 \to @_m\Box(\psi_2 \to \#))$, then $L(\varphi)$ is $\psi_0 \to @_n\Box(\psi_1 \to @_m\Box(\psi_2 \to \varphi))$. Intuitively, this notion allows us to capture the unique path from a label in a tree of a tree sequent to the root label of the tree.

Table 2 presents our Hilbert system $\mathsf{H}\mathbf{EFL}$. The underlying idea of the system is: on the top of the propositional part (`Taut` and `MP`), we combine the axiomatization of modal logic $\mathbf{K}$ for the modal operator $\Box$ and the axiomatization of a basic hybrid logic $\mathbf{K}_{\mathcal{H}(@)}$ for the modal operator F, with some modification (we need to modify $\mathbf{BG}$, the rule of *bounded generalization*, with the help of necessity forms), and then we add three interaction axioms: ($\mathtt{Rigid}_{=}$), ($\mathtt{Rigid}_{\neq}$), and ($\mathtt{DCom@\Box}$). We note that the axiom ($\mathtt{DCom@\Box}$) is also used for axiomatizing the *dependent product* of two hybrid logics in [17]. Let us define the notion of provability in $\mathsf{H}\mathbf{EFL}$ in as usual. We write $\vdash_{\mathsf{HEFL}} \varphi$ to means that φ is provable in $\mathsf{H}\mathbf{EFL}$.

Table 2. Hilbert System $\mathsf{H}\mathbf{EFL}$

(Taut) all propositional tautologies	(MP) From φ and $\varphi \to \psi$, infer ψ
($\mathtt{K}_\Box$) $\Box(p \to q) \to (\Box p \to \Box q)$	($\mathtt{Nec}_\Box$) From φ, infer $\Box\varphi$
($\mathtt{K}_\mathsf{F}$) $\mathsf{F}(p \to q) \to (\mathsf{F}p \to \mathsf{F}q)$	($\mathtt{Nec}_\mathsf{F}$) From φ, infer $\mathsf{F}\varphi$
($\mathtt{K}_@$) $@_n(p \to q) \to (@_n p \to @_n q)$	(Ref) $@_n n$
(Selfdual) $\neg @_n p \leftrightarrow @_n \neg p$	(Elim) $@_n p \to (n \to p)$
(Agree) $@_n @_m p \to @_m p$	(Back) $@_n p \to \mathsf{F}@_n p$
($\mathtt{Rigid}_=$) $@_n m \to \Box @_n m$	($\mathtt{Rigid}_\neq$) $\neg @_n m \to \Box \neg @_n m$
(DCom@$\Box$) $@_n \Box @_n p \leftrightarrow @_n \Box p$	
(US) From φ, infer $\varphi\sigma$, where σ is a uniform substitution.	
(Name) From $n \to \varphi$, infer φ, where n is fresh in φ.	
(L(BG)) From $L(@_n\langle\mathsf{F}\rangle m \to @_m\varphi)$, infer $L(@_n\mathsf{F}\varphi)$, where m is fresh in $L(@_n\mathsf{F}\varphi)$.	

Proposition 2. *All the following are provable in* $\mathsf{H}\mathbf{EFL}$.

1. $@_m @_n \varphi \leftrightarrow @_n \varphi$.
2. $n \to (@_n \varphi \leftrightarrow \varphi)$.
3. $@_n m \to (@_n \varphi \leftrightarrow @_m \varphi)$.
4. $@_n m \leftrightarrow @_m n$.
5. $@_n(\varphi \to \psi) \leftrightarrow (@_n \varphi \to @_n \psi)$.
6. $@_n m \to (\varphi[n/k] \leftrightarrow \varphi[m/k])$.

The following translation is a key to specify our Hilbert system $\mathsf{H}\mathbf{EFL}$.

Definition 3 (formulaic translation). *Given a set Θ of labelled formulas and a label α, we define $\Theta_\alpha := \{\, \varphi \mid \alpha : \varphi \in \Theta \,\}$. Let $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ be a tree sequent. Then the formulaic translation of the sequent at α is defined as:*

$$\left[\!\!\left[\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta \right]\!\!\right]_\alpha := \bigwedge \Gamma_\alpha \to \bigvee \left(\Delta_\alpha, @_{n_1}\Box \left[\!\!\left[\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta \right]\!\!\right]_{\beta_1}, \ldots, @_{n_k}\Box \left[\!\!\left[\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta \right]\!\!\right]_{\beta_k} \right),$$

where β_i is an n_i-child of α, β_is enumerate all children of α, $\bigwedge \emptyset := \top$, and $\bigvee \emptyset := \bot$.

The formulaic translation of a tree sequent of Fig. 1 of Section 3 at the root 0 is

$$@_n \varphi \to (@_n \psi \vee @_n \Box(\top \to @_k \theta) \vee @_k \Box(@_m \rho \to \bot)).$$

Theorem 3. *If a tree sequent $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ is provable in $\mathsf{T}\mathbf{EFL}$ then the formulaic translation $[\![\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta]\!]_i$ is provable in $\mathsf{H}\mathbf{EFL}$, where a natural number i is the root of $\mathcal{T}$.*

Proof. By induction on height n of a derivation of $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ in $\mathsf{T}\mathbf{EFL}$, where i is the root of the tree $\mathcal{T}$. We skip the base case where $n = 0$. Let $n > 0$. It is remarked that, when the sequent is obtained by (rep_l), $(\mathsf{ref}_=)$, $(@L)$, or $(@R)$, respectively, the translation of the sequent at the root is provable by Proposition 2 (6), the axiom (Ref), (Agree), or Proposition 2 (1), respectively. Here we focus on the case where $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ is obtained by $(\mathsf{rigid}_=)$, $(\mathsf{F}R)$, or $(\Box R)$, since these are the cases where we need to be careful and the other cases are easy to establish.

($\Box L$) Suppose that $\alpha : @_n\Box\varphi, \Gamma' \overset{\mathcal{T}}{\Rightarrow} \Delta$ is obtained by ($\Box L$) from $\beta : @_n\varphi, \Gamma' \overset{\mathcal{T}}{\Rightarrow} \Delta$, where $\beta \in \mathcal{T}$ is an n-child of α. By induction hypothesis, we obtain $\vdash_{\mathbf{HEFL}} \left[\!\left[\beta : @_n\varphi, \Gamma' \overset{\mathcal{T}}{\Rightarrow} \Delta\right]\!\right]_i$. We show that $\vdash_{\mathbf{HEFL}} \left[\!\left[\alpha : @_n\Box\varphi, \Gamma' \overset{\mathcal{T}}{\Rightarrow} \Delta\right]\!\right]_i$. Let $(\alpha_0, \alpha_1, \ldots, \alpha_l)$ be the unique path from α ($\equiv \alpha_l$) to the root i ($\equiv \alpha_0$) of tree $\mathcal{T}$. By induction on $0 \leqslant h \leqslant l$, we show that $\vdash_{\mathbf{HEFL}} \left[\!\left[\beta : @_n\varphi, \Gamma' \overset{\mathcal{T}}{\Rightarrow} \Delta\right]\!\right]_{\alpha_{l-h}} \rightarrow \left[\!\left[\alpha : @_n\Box\varphi, \Gamma' \overset{\mathcal{T}}{\Rightarrow} \Delta\right]\!\right]_{\alpha_{l-h}}$.
Let $h = 0$ and so $\alpha_{l-h} = \alpha$. It suffices to show that a formula of the form

$$(\gamma_1 \rightarrow (\delta \vee @_n\Box((\gamma_2 \wedge @_n\varphi) \rightarrow \psi_2))) \rightarrow ((@_n\Box\varphi \wedge \gamma_1) \rightarrow (\psi_1 \vee @_n\Box(\gamma_2 \rightarrow \psi_2)))\,.$$

is provable in HEFL. This reduces to the provability of

$$@_n\Box\varphi \wedge @_n\Box((\gamma_2 \wedge @_n\varphi) \rightarrow \psi_2)) \rightarrow @_n\Box(\gamma_2 \rightarrow \psi_2))$$

in HEFL. This holds by the axiom ($\mathtt{Dcom}\Box@$) $@_n\Box@_n\varphi \leftrightarrow @_n\Box\varphi$.
Let $h > 0$. But this case is shown with the help of ($\mathsf{Nec}_\Box$) and ($\mathsf{Nec}_@$). This completes our induction on h. So we conclude $\vdash_{\mathbf{HEFL}} \left[\!\left[\alpha : @_n\Box\varphi, \Gamma' \overset{\mathcal{T}}{\Rightarrow} \Delta\right]\!\right]_i$.

($\mathsf{F}R$) Suppose that $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta', \alpha : @_n\mathsf{F}\varphi$ is obtained by ($\mathsf{F}R$) from $\alpha : @_n\langle\mathsf{F}\rangle m, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta', \alpha : @_m\varphi$ where m is fresh in the conclusion. By induction hypothesis, we have $\vdash_{\mathbf{HEFL}} \left[\!\left[\alpha : @_n\langle\mathsf{F}\rangle m, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta', \alpha : @_m\varphi\right]\!\right]_i$, which is equivalent to $\vdash_{\mathbf{HEFL}} L(@_n\langle\mathsf{F}\rangle m \rightarrow @_m\varphi)$ for some necessitation form L. Fix such necessitation form L. By the inference rule $L(\mathtt{BG})$ of HEFL, we can obtain $\vdash_{\mathbf{HEFL}} L(@_n\mathsf{F}\varphi)$, which is equivalent to $\vdash_{\mathbf{HEFL}} \left[\!\left[\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta', \alpha : @_n\mathsf{F}\varphi\right]\!\right]_i$.

($\mathsf{rigid}_=$) Instead of dealing with a general case, we handle a simple example of $\mathcal{T}$ to extract an essence of this case, where we need to use the axioms ($\mathtt{Rigid}_=$) and ($\mathtt{Rigid}_\neq$). Let $\mathcal{T}$ consists of three labels, i.e., the root i, a k-child α of i and a k'-child β of i. Let us suppose that $\beta : @_n m, \Gamma' \overset{\mathcal{T}}{\Rightarrow} \Delta$ is obtained by ($\mathsf{rigid}_=$) from $\alpha : @_n m, \Gamma' \overset{\mathcal{T}}{\Rightarrow} \Delta$. In what follows, for every $\eta \in \mathcal{T}$, let us write $\bigwedge \Gamma'_\eta$ and $\bigvee \Delta_\eta$ by γ_η and δ_η, respectively. Here we note that the following hold:

(β to i) $\vdash_{\mathbf{HEFL}} (@_n m \wedge @_{k'}\Box(@_n m \wedge \gamma_\beta \rightarrow \delta_\beta)) \rightarrow @_{k'}\Box(\gamma_\beta \rightarrow \delta_\beta)$

(i to α) $\vdash_{\mathbf{HEFL}} (\neg@_n m \vee @_k\Box(\gamma_\alpha \rightarrow \delta_\alpha)) \rightarrow @_k\Box((@_n m \wedge \gamma_\alpha) \rightarrow \delta_\alpha)$.

For (β to i), it suffices to show $\vdash_{\mathbf{HEFL}} @_n m \rightarrow @_k\Box@_n m$ by ($\mathtt{Rigid}_=$), which holds by the distribution of $@$ over the implication and Proposition 2 (1). For (i to α), it suffices to show $\vdash_{\mathbf{HEFL}} \neg@_n m \rightarrow @_k\Box\neg@_n m$, which holds by ($\mathtt{Rigid}_\neq$).
By induction hypothesis, we obtain $\vdash_{\mathbf{HEFL}} \left[\!\left[\alpha : @_n m, \Gamma' \overset{\mathcal{T}}{\Rightarrow} \Delta\right]\!\right]_i$, i.e.,

$$\vdash_{\mathbf{HEFL}} \gamma_i \rightarrow (\delta_i \vee @_k\Box(\gamma_\alpha \rightarrow \delta_\alpha) \vee @_{k'}\Box((@_n m \wedge \gamma_\alpha) \rightarrow \delta_\alpha))\,.$$

It follows from item (β to i) that

$$\vdash_{\mathbf{HEFL}} (@_n m \wedge \gamma_i) \rightarrow (\delta_i \vee @_k\Box(\gamma_\alpha \rightarrow \delta_\alpha) \vee @_{k'}\Box(\gamma_\alpha \rightarrow \delta_\alpha))\,.$$

By this and item (i to α), we can establish:

$$\vdash_{\mathbf{HEFL}} \gamma_i \rightarrow (\delta_i \vee @_k\Box((@_n m \wedge \gamma_\alpha) \rightarrow \delta_\alpha) \vee @_{k'}\Box(\gamma_\alpha \rightarrow \delta_\alpha))\,,$$

which is equivalent to: $\vdash_{\mathbf{HEFL}} \left[\!\left[\beta : @_n m, \Gamma' \overset{\mathcal{T}}{\Rightarrow} \Delta\right]\!\right]_i$, as desired. □

In what follows in this section, we prove the soundness of $\mathsf{H}\mathbf{EFL}$ for the tree sequent calculus $\mathsf{T}\mathbf{EFL}$ with cut rule. The cut rule is necessary to prove the following.

Lemma 3. *The rules $(\to R)$, $(\Box R)$, $(@R)$, and $(@L)$ are invertible, i.e., if the lower sequent is provable in $\mathsf{T}\mathbf{EFL}$ then the upper sequent is also provable in $\mathsf{T}\mathbf{EFL}$.*

Theorem 4. *If φ is provable in $\mathsf{H}\mathbf{EFL}$, then $\overset{\mathcal{T}}{\Rightarrow} \alpha : @_n\varphi$ is provable in $\mathsf{T}\mathbf{EFL}$ for all fresh nominals n in φ, tree $\mathcal{T}$ and $\alpha \in \mathcal{T}$.*

Proof. Suppose that there is a proof $(\varphi_0, \dots, \varphi_h)$ of φ in $\mathsf{H}\mathbf{EFL}$. By induction on $0 \leqslant j \leqslant h$, we show that $\overset{\mathcal{T}}{\Rightarrow} \alpha : @_n\varphi_j$ is provable in $\mathsf{T}\mathbf{EFL}$ for all fresh nominals n in φ_j and $\alpha \in \mathcal{T}$. Since the space is limited, we demonstrate some cases. Let us start with ($\mathtt{Rigid}_{=}$), which is shown by the left derivation below. Now we move to ($\mathtt{DCom}@\Box$). We show the right-to-left direction alone, since the converse direction is shown similarly. Let us see the right derivation below, from which we can obtain the provability of $\overset{\mathcal{T}}{\Rightarrow} \alpha : @_m(@_n\Box@_n p \to @_n\Box p)$ in $\mathsf{T}\mathbf{EFL}$. Now we deal with some of the inference rules below.

$$\dfrac{\dfrac{\dfrac{\dfrac{\alpha \cdot_k i : @_n m \overset{\mathcal{T} \cup \{\alpha \cdot_k i\}}{\Rightarrow} \alpha \cdot_k i : @_n m}{\alpha : @_n m \overset{\mathcal{T} \cup \{\alpha \cdot_k i\}}{\Rightarrow} \alpha \cdot_k i : @_n m}\,(@R)}{\alpha : @_n m \overset{\mathcal{T} \cup \{\alpha \cdot_k i\}}{\Rightarrow} \alpha \cdot_k i : @_k @_n m}\,(\Box R)}{\alpha : @_n m \overset{\mathcal{T}}{\Rightarrow} \alpha : @_k \Box @_n m}\,(@L)}{\alpha : @_k @_n m \overset{\mathcal{T}}{\Rightarrow} \alpha : @_k \Box @_n m}\,(\to R)}{\overset{\mathcal{T}}{\Rightarrow} \alpha : @_k(@_n m \to \Box @_n m)}$$

$$\dfrac{\dfrac{\dfrac{\alpha \cdot_n i : @_n p \overset{\mathcal{T} \cup \{\alpha \cdot_n i\}}{\Rightarrow} \alpha \cdot_n i : @_n p}{\alpha \cdot_n i : @_n @_n p \overset{\mathcal{T} \cup \{\alpha \cdot_n i\}}{\Rightarrow} \alpha \cdot_n i : @_n p}\,(\Box L)}{\alpha : @_n \Box @_n p \overset{\mathcal{T} \cup \{\alpha \cdot_n i\}}{\Rightarrow} \alpha : \cdot_n i : @_n p}\,(\Box R)}{@_n \Box @_n p \overset{\mathcal{T}}{\Rightarrow} \alpha : @_n \Box p}$$

($L(\mathtt{BG})$) Let $\varphi_j \equiv \Box\psi$ be obtained by $(L(\mathtt{BG}))$. For simplicity, we assume that $L \equiv @_n\Box(\psi \to \#)$ here, while our argument is easily extended to a general case. Fix any tree $\mathcal{T}$, $\alpha \in \mathcal{T}$ and fresh nominal m. By induction hypothesis, $\overset{\mathcal{T}}{\Rightarrow} @_n\Box(\psi \to (@_n\langle \mathsf{F}\rangle m \to @_m\varphi))$ is provable in $\mathsf{T}\mathbf{EFL}$, where m satisfies the freshness condition. By applying Lemma 3 repeatedly, we obtain the provability of $\alpha \cdot_n i : @_n\psi, \alpha \cdot_n i : @_n\langle \mathsf{F}\rangle m \overset{\mathcal{T} \cup \{\alpha \cdot_n i\}}{\Rightarrow} \alpha \cdot_n i : @_m\varphi$ in $\mathsf{T}\mathbf{EFL}$ for some fresh i. Then we proceed as follows:

$$\dfrac{\dfrac{\dfrac{\dfrac{\alpha \cdot_n i : @_n\psi, \alpha \cdot_n i : @_n\langle \mathsf{F}\rangle m \overset{\mathcal{T} \cup \{\alpha \cdot_n i\}}{\Rightarrow} \alpha \cdot_n i : @_m\varphi}{\alpha \cdot_n i : @_n\psi \overset{\mathcal{T} \cup \{\alpha \cdot_n i\}}{\Rightarrow} \alpha \cdot_n i : @_n \mathsf{F}\varphi}\,(\mathsf{F}R)}{\alpha \cdot_n i : @_n\psi \overset{\mathcal{T} \cup \{\alpha \cdot_n i\}}{\Rightarrow} \alpha \cdot_n i : @_n @_n \mathsf{F}\varphi}\,(@R)}{\overset{\mathcal{T} \cup \{\alpha \cdot_n i\}}{\Rightarrow} \alpha \cdot_n i : @_n(\psi \to @_n \mathsf{F}\varphi)}\,(\to L)}{\overset{\mathcal{T}}{\Rightarrow} \alpha : @_n\Box(\psi \to @_n \mathsf{F}\varphi)}\,(\Box R)\quad,$$

as requred. □

Corollary 2 (Soudness and Completenss of $\mathsf{H}\mathbf{EFL}$). *The following are all equivalent: for every formula φ,*

1. *φ is valid in the class of all models,*
2. *$\overset{\mathcal{T}}{\Rightarrow} \alpha : @_n\varphi$ is provable in $\mathsf{T}\mathbf{EFL}^-$ for all fresh nominals n in φ, tree $\mathcal{T}$ and $\alpha \in \mathcal{T}$,*
3. *$\overset{\mathcal{T}}{\Rightarrow} \alpha : @_n\varphi$ is provable in $\mathsf{T}\mathbf{EFL}$ for all fresh nominals n in φ, tree $\mathcal{T}$ and $\alpha \in \mathcal{T}$,*
4. *φ is provable in $\mathsf{H}\mathbf{EFL}$.*

Proof. Item 1 is equivalent to the following: $\overset{\mathcal{T}}{\Rightarrow} \alpha : @_n\varphi$ is true for all pairs $(\mathfrak{M}, f)$ of models and assignments, fresh nominals n in φ, tree $\mathcal{T}$, and $\alpha \in \mathcal{T}$. Then the equivalence between items 1, 2 and 3 hold by Corollary 1. The direction from item 4 to item 3 holds by Theorem 4. Finally, the direction from item 3 to item 4 is established as follows. Suppose item 3. Let n be a fresh nominal α. By the supposition, $\overset{\{0\}}{\Rightarrow} 0 : @_n\varphi$ is provable in $\mathsf{T}\mathbf{EFL}$. It follows from Theorem 3 that $\vdash_{\mathbf{HEFL}} [\![\overset{\{0\}}{\Rightarrow} 0 : @_n\varphi]\!]_0$, which implies $\vdash_{\mathbf{HEFL}} @_n\varphi$. By the axiom (Elim), we obtain $\vdash_{\mathbf{HEFL}} n \to \varphi$ hence $\vdash_{\mathbf{HEFL}} \varphi$ by (Name), as required. □

6 Extensions of Epistemic Logic of Friendship

This section outlines how we extend our tree sequent calculus $\mathsf{T}\mathbf{EFL}$ and Hibert system $\mathsf{H}\mathbf{EFL}$. In particular, we discuss extensions where $\Box$ follows $\mathbf{S4}$ or $\mathbf{S5}$ axioms and/or the friendship relation $\asymp_w$ satisfies some universal properties such as irreflexivity, symmetry, etc. ($w \in W$).

Let us denote a set $\{\, \Box p \to p, \Box p \to \Box\Box p \,\}$ by $\mathbf{S4}$ and a set $\mathbf{S4} \cup \{\, p \to \Box\neg\Box\neg p \,\}$ by $\mathbf{S5}$. Let us consider formulas of the form $@_n m$ or $@_n\langle \mathsf{F}\rangle m$, which are denoted by ρ_i, ρ_{ij}, etc. below. Let us consider a formula φ of the following form:

$$(\rho_1 \wedge \cdots \wedge \rho_h) \to \bigvee_{1 \le j \le l} \left(\rho_{j1} \wedge \cdots \wedge \rho_{jk_j}\right),$$

where we note that h and l are possibly zero. We say that a formula of such form is a *u-geometric implication*. The name of u-geometric implication is coming from the notion of geometric formula (see, e.g., [6] in the context of hybrid logic) and, unlike geometric formulas, we do not consider the existential quantification to the conclusion of the implication. This is just for simplicity. The corresponding frame property of a u-geometric implication is obtained by regarding $@_n m$ or $@_n\langle \mathsf{F}\rangle m$ by "$a_n = a_m$" and "$a_n \asymp_w a_m$" and putting the universal quantifiers for all agents and w. For example, irreflexivity and symmetry of $\asymp_w$ are defined by $@_n\langle \mathsf{F}\rangle n \to \bot$ and $@_n\langle \mathsf{F}\rangle m \to @_m\langle \mathsf{F}\rangle n$, respectively. When Λ is one of $\mathbf{S4}$ and $\mathbf{S5}$ and Θ is a finite set of u-geometric implications, a Hilbert system $\mathsf{H}\mathbf{EFL}(\Lambda \cup \Theta)$ is defined as the axiomatic extension of $\mathsf{H}\mathbf{EFL}$ by new axioms $\Lambda \cup \Theta$.

Now let us move to tree sequent systems. First, we introduce an inference rule for a u-geometric implication. For a u-geometric implication φ displayed above, we can define the corresponding inference rule $(\mathsf{ugi}(\varphi))$ for tree sequent calculus as follows (cf. [6]):

$$\frac{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta, \alpha : \rho_1 \quad \ldots \quad \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta, \alpha : \rho_h \quad \alpha : \rho_{11}, \ldots, \alpha : \rho_{1k_1}, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta \quad \ldots \quad \alpha : \rho_{l1}, \ldots, \alpha : \rho_{lk_l}, \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}\ (\mathsf{ugi}(\varphi))$$

When $\asymp_w$ is irreflexive or symmetric for all $w \in W$, we can obtain the following rule $(\mathsf{irr}_{\asymp})$ or $(\mathsf{sym}_{\asymp})$, respectively:

$$\frac{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta, \alpha : @_n\langle \mathsf{F}\rangle n}{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}\ (\mathsf{irr}_{\asymp}) \qquad \frac{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta, \alpha : @_n\langle \mathsf{F}\rangle m \quad \alpha : @_m\langle \mathsf{F}\rangle n\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}{\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta}\ (\mathsf{sym}_{\asymp}).$$

Let Λ be one of $\mathbf{S4}$ and $\mathbf{S5}$ and Θ be a possibly empty finite set of u-geometric implications. In what follows, we define the tree sequent system $\mathsf{T}\mathbf{EFL}(\Lambda;\Theta)$. Recall that the side condition $\ddagger$ of the rule $(\Box L)$ of Table 1. Depending on the choice of Λ, we change the side condition $\ddagger$ into the following one in $\mathsf{T}\mathbf{EFL}(\Lambda;\Theta)$:

- $\ddagger_{\mathbf{S4}}$: $\alpha \preceq_n \beta$, where $\preceq_n$ is the reflexive transitive closure of the n-children relation.
- $\ddagger_{\mathbf{S5}}$: $\alpha \sim_n \beta$, where $\sim_n$ is the reflexive, symmetric, transitive closure of the n-children relation.

Second, we extend the system with a set $\{\, (\mathsf{ugi}(\varphi)) \mid \varphi \in \Theta \,\}$ of inference rules to finish to define the system $\mathsf{T}\mathbf{EFL}(\Lambda;\Theta)$. We define $\mathsf{T}\mathbf{EFL}(\Lambda;\Theta)^-$ as the system $\mathsf{T}\mathbf{EFL}(\Lambda;\Theta)$ without the cut rule.

Given a set Ψ of formulas and a frame $\mathfrak{F} = (W, A, (R_a)_{a\in A}, (\asymp_w)_{w\in W})$ (a model without a valuation), we say that Ψ is *valid* in $\mathfrak{F}$ (notation: $\mathfrak{F} \models \Psi$) if $(\mathfrak{F}, V), (w, a) \models \psi$ for all $\psi \in \Psi$, valuations V and pairs $(w, a) \in W \times A$. We define a class $\mathbb{M}_\Psi$ of models as $\{\, (\mathfrak{F}, V) \mid \mathfrak{F} \models \Psi \,\}$. While we omit the detail of the proof, we can obtain the following two theorems by similar arguments to $\mathsf{T}\mathbf{EFL}$ and $\mathsf{H}\mathbf{EFL}$.

Theorem 5. *Let Λ be one of $\mathbf{S4}$ and $\mathbf{S5}$ and Θ be a possibly empty finite set of u-geometric implications. The following are all equivalent:*

1. $\mathfrak{M}, f \models \Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ *for all models* $\mathfrak{M} \in \mathbb{M}_{\Lambda\cup\Theta}$ *and all assignments* f.
2. $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ *is provable in* $\mathsf{T}\mathbf{EFL}(\Lambda;\Theta)^-$.
3. $\Gamma \overset{\mathcal{T}}{\Rightarrow} \Delta$ *is provable in* $\mathsf{T}\mathbf{EFL}(\Lambda;\Theta)$.

Therefore, $\mathsf{T}\mathbf{EFL}(\Lambda;\Theta)$ enjoys the cut-elimination theorem. Moreover, for every formula φ, φ is valid in $\mathbb{M}_{\Lambda\cup\Theta}$ iff φ is provable in $\mathsf{H}\mathbf{EFL}(\Lambda \cup \Theta)$.

7 Further Directions

This paper positively answered the question if the set of all valid formulas of $\mathbf{EFL}$ in the class of all models is axiomatizable. We list some directions for further research.

1. Is $\mathsf{H}\mathbf{EFL}$ or $\mathsf{T}\mathbf{EFL}$ decidable?
2. Is it possible to provide a syntactic proof of the cut elimination theorem of $\mathsf{T}\mathbf{EFL}$?
3. Can we reformulate our sequent calculus into a G3-style calculus, i.e., a contraction fee calculus, all of whose rule are height-preserving invertible?
4. Provide a G3-style labelled sequent calculus for $\mathbf{EFL}$ based on the idea of doubly labelled formula $(x, y) : \varphi$. This is an extension of G3-style labelled sequent calculus for modal logic in [15,14].
5. Prove the semantic completeness of $\mathsf{H}\mathbf{EFL}$ and its extensions by specifying the notion of *canonical model*.
6. Can we apply our technique of this paper obtain a Hilbert-system of *Term Modal Logics* which is proposed in [8]?

References

1. C. Areces and B. ten Cate. Hybrid logics. In Patrick Blackburn, Johan van Benthem, and Frank Wolter, editors, *Handbook of Modal Logic*, pages 821–868. Elsevier, 2007.
2. A. Baltag, L. Moss, and S. Solecki. The logic of public announcements, common knowledge and private suspicions. In *Proceedings of TARK*, pages 43–56, Los Altos, 1989. Morgan Kaufmann Publishers.
3. P. Blackburn. Arthur Prior and hybrid logic. *Synthese*, 150(3):329–372, 2006.
4. P. Blackburn and M. Tzakova. Hybrid completeness. *Logic Journal of IGPL*, 6(4):625–650, 1998.
5. T. Braüner. *Hybrid Logic and Its Proof-Theory*, volume 37 of *Applied Logic Series*. Springer, 2011.
6. K. Brünnler. Deep sequent systems for modal logic. *Archive for Mathematical Logic*, 48:551577, 2009.
7. Z. Christoff, J. U. Hansen, and C. Proetti. Reflecting on social influence in networks. *Journal of Logic, Language and Information*, 25(3):299–333, 2016.
8. M. Fitting, L. Thalmann, and A. Voronkov. Term-modal logics. *Studia Logica*, 69(1):133–169, 2001.
9. G. Gargov, S. Passy, and T. Tinchev. Modal environment for Boolean speculations (preliminary report). In D. Skordev, editor, *Mathematical Logic and its Applications. Proceedings of the Summer School and Conference dedicated to the 80th Anniversary of Kurt Gödel*, pages 253–263. Plenum Press, Druzhba, 1987.
10. P. Girard, J. Seligman, and F. Liu. General dynamic dynamic logic. In S. Ghilardi T. Bolander, T Braüner and L. S. Moss, editors, *Advances in Modal Logics*, volume 9, pages 239–260. College Publications, 2012.
11. R. Goldblatt. *Axiomatizing the logic of computer programming*, volume 130 of *Lecture Notes in Computer Science*. Springer-Verlag, 1982.
12. J. Hintikka. *Knowledge and Belief: An Introduction to the Logic of the Two Notions*. Cornell University Press, Cornell, 1962.
13. R. Kashima. Cut-free sequent calculi for some tense logics. *Studia Logica*, 53:119–135, 1994.
14. S. Negri. Proof analysis in modal logic. *Journal of Philosophical Logic*, 34:507544, 2005.
15. S. Negri and J. Von Plato. *Proof Analysis*. Cambridge University Press, 2011.
16. J. A. Plaza. Logics of public communications. In M. L. Emrich, M. S. Pfeifer, M Hadzikadic, and Z. W. Ras, editors, *Proceedings of the 4th International Symposium on Methodologies for Intelligent Systems*, pages 201–216, 1989.
17. K. Sano. Axiomatizing hybrid products: How can we reason many-dimensionally in hybrid logic? *Journal of Applied Logic*, 8(4):459–474, December 2010.
18. J. Seligman. Internalization: The case of hybrid logics. *Journal of Logic and Computation*, 11(5):671–689, 2001.
19. J. Seligman, F. Liu, and P. Girard. Logic in the community. In *Proceedings of the 4th Indian Conference on Logic and its Applications*, volume 6521 of *Lecture Notes in Computer Science*, pages 178–188, 2011.
20. J. Seligman, F. Liu, and P. Girard. Knowledge, friendship and social announcement. In Johan van Benthem and Fenrong Liu, editors, *Logic Across the University: Foundations and Applications*, volume 47 of *Studies in Logic*, pages 445–469. College Publications, 2013.